RESEARCH PAPER

ABSTRACT

Superficial white matter damage in anti-NMDA receptor encephalitis

Owen Robert Phillips,¹ Shantanu H Joshi,² Katherine L Narr,² David W Shattuck,² Manpreet Singh,¹ Margherita Di Paola,^{3,4} Christoph J Ploner,⁵ Harald Prüss,^{5,6} Friedemann Paul,^{5,7,8,9} Carsten Finke^{5,10}

► Additional material is published online only. To view please visit the journal online (http://dx.doi.org/10.1136/ jnnp-2017-316822).

For numbered affiliations see end of article.

Correspondence to

Professor Carsten Finke, Department of Neurology, Charité–Universitätsmedizin Berlin, Charitéplatz 1, 10117 Berlin, Germany; carsten. finke@charite.de and Professor Friedemann Paul, Department of Neurology Charité University Medicine Berlin, Berlin, Germany; friedemann.paul@ charite.de

Received 6 July 2017 Revised 9 October 2017 Accepted 19 October 2017 Published Online First 3 November 2017 **Background** Clinical brain MRI is normal in the majority of patients with anti-*N*-methyl-D-aspartate receptor (NMDAR) encephalitis. However, extensive deep white matter damage wasrecently identified in these patients using diffusion weighted imaging. Here, our aim was to study a particularly vulnerable brain compartment, the late myelinating superficial white matter.

Methods Forty-six patients with anti-NMDAR encephalitis were included. Ten out of these were considered neurologically recovered (modified Rankin scale of zero), while 36 patients were non-recovered. In addition, 30 healthy controls were studied. MRI data were collected from all subjects and superficial white matter mean diffusivity derived from diffusion tensor imaging was compared between groups in whole brain, lobar and vertex-based analyses. Patients underwent comprehensive cognitive testing, and correlation analyses were performed between cognitive performance and superficial white matter integrity.

Results Non-recovered patients showed widespread superficial white matter damage in comparison to recovered patients and healthy controls. Vertex-based analyses revealed that damage predominated in frontal and temporal lobes. In contrast, the superficial white matter was intact in recovered patients. Importantly, persistent cognitive impairments in working memory, verbal memory, visuospatial memory and attention significantly correlated with damage of the superficial white matter in patients.

Conclusions Anti-NMDAR encephalitis is associated with extensive superficial white matter damage in patients with incomplete recovery. The strong association with impairment in several cognitive domains highlights the clinical relevance of white matter damage in this disorder and warrants investigations of the underlying pathophysiological mechanisms.

INTRODUCTION

Anti-N-methyl-D-aspartate receptor (NMDAR) encephalitis is an autoimmune encephalitis that exhibits a characteristic neuropsychiatric syndrome and has a severe and prolonged clinical course.¹ Clinical routine MRI is normal in most patients, but advanced imaging analyses, including resting state functional MRI (fMRI), volumetric analyses and diffusion tensor imaging, have identified structural and functional imaging correlates of the disease.^{2–4} Recently, we observed a characteristic pattern of whole-brain functional connectivity alterations that correlated with memory impairment and psychiatric symptoms in anti-NMDAR encephalitis.⁵ Moreover, extensive white matter changes within the large deep white matter fibres that were shown to correlate with disease severity were described.² However, the exact contribution of white matter damage to disease pathophysiology remains elusive. To address this issue, we examined the integrity of the short-range association fibres (U-fibres) and intracortical myelin, which is made up largely of oligodendrocytes, at the interface between cortical grey matter and deep white matter (here called the superficial white matter; see table 1 for a brief overview) to test links between structural and functional impairment in NMDAR encephalitis. Several lines of evidence support the hypothesis of disease-related damage to the superficial white matter and why it may be linked to functional impairments. First, the superficial white matter is a highly vulnerable area that matures after the deep white matter.⁶⁻⁸ Moreover, the oligodendrocytes in the superficial white matter myelinate axon segments with fewer wraps than in the deep white matter,⁹ thus rendering axons more susceptible to impairments.¹⁰ In addition, the superficial white matter has been shown to be especially vulnerable in schizophrenia,¹¹ a disease which shares pathophysiological mechanisms (NMDAR dysfunction), clinical symptoms (including positive and negative symptoms as well as cognitive deficits) and neuroimaging findings (hippocampal atrophy, damage of deep white matter, altered functional connectivity within the frontoparietal control network and the ventral attention network) with anti-NMDAR encephalitis.²³

Importantly, although patients with anti-NMDAR encephalitis frequently suffer from a severe clinical course, many patients respond well to treatment and return to a relatively high level of function.^{12 13} Together with recent neuroimaging findings, this suggests that functional impairment and structural damage of the brain in patients with active symptoms may recover substantially. Given its vulnerability, we investigated the superficial white matter with the hypothesis that it may be adversely affected in patients with ongoing disease activity, but may recover in patients who no longer exhibit symptoms. Furthermore, we investigated a possible relationship between integrity of superficial white matter and cognitive impairment.

To cite: Phillips OR, Joshi SH, Narr KL, et al. J Neurol Neurosurg Psychiatry 2018;89:518–525.



Table 1 Differences between superficial and deep white matter						
SWM	Deep white matter					
Short fibres	Long fibres					
Small diameter fibres	Larger diameter fibres					
Intracortical connections	Intrahemispheric and interhemispheric connections					
Late myelinating	Early myelinating compared with SWM					
Oligodendrocytes wrap many axons segments	Oligodendrocytes wrap few axon segments					
Less myelin wraps around the axon	More myelin wraps around the axon					
Complex arrangement	Less complex arrangement compared with SWM					

SWM, superficial white matter.

METHODS

Forty-six patients with anti-NMDAR encephalitis were recruited from the Department of Neurology at Charité - Universitätsmedizin Berlin (age: mean±SD: 26.67±8.45 years, 40F/6M). All patients had a characteristic clinical presentation¹⁴ and were tested positive for IgG NMDAR antibodies using an immunofluorescence cell-based assay expressing recombinant NMDAR. In addition, all patients tested negative for AMPAR, GABA-B-R, LGI1 und CASPR2 antibodies. Similarly, MOG and AQP4 antibodies were not detected (tested in 21/46 patients). The control group comprised 30 healthy subjects without history of neurological or psychiatric disorders (age: 27.53±8.42 years, 24F/6M). Resting state fMRI, structural MRI and optical coherence tomography analyses of some patients have previously been reported.^{2 3 15} Patients were studied 27.6 ± 4.6 months after symptom onset and 12.1 ± 2.4 months after the end of the last acute care hospitalisation. The patient's disease severity was independently assessed at the time of MRI studies by two experienced neurologists using the modified Rankin scale (mRS). Ten patients with an mRS score of 0 were considered clinically recovered (recovered patients, age, 26.60 ± 7.38 years, 8F/2M), while 36 patients had residual deficits as indicated by mRS scores >0(non-recovered patients, age, 26.69±8.83 years, 32F/4M; mRS=1, 15 patients; mRS=2, 16 patients; mRS=3, 6 patients). Importantly, the mRS is a general measure of function and does not capture subtle deficits, including mild to moderate cognitive dysfunction, which may still be present. In addition to the mRS at the time of study, the maximal mRS during the acute disease phase was assessed (table 2). All study participants gave written informed consent for research and publication.

Cognitive assessment

All study participants underwent cognitive testing with a comprehensive test battery that covered attention (alertness and divided attention), verbal and non-verbal short-term and working memory (digit span forward and backward, block tapping forward and backward), verbal and visuospatial memory (Rey Auditory Verbal Learning Test (RAVLT) and Rey-Osterrieth Complex Figure Test (ROCF)) and executive function (computerised go/no-go test), as described in detail previously.¹⁶

MRI data acquisition

MRI data were acquired on a Siemens Magnetom Tim Trio 3T scanner equipped with a 12-channel phased-array head coil and included the following sequences: (i) 3D 1 mm isotropic magnetisation-prepared rapid gradient echo sequence (repetition time(TR)=1900 ms, echo time (TE)=2.55 ms, TI=900ms, flip angle=9°, field-of-view (FOV)= 240×240 mm², matrix size= 240×240 , 176 slices, slice thickness=1 mm) for structural acquisition and (ii) a single-shot echo-planar imaging sequence for diffusion MRI acquisition (TR=7500 ms, TE=86 ms, FOV= 240×240 mm², voxel size= $2.5 \times 2.5 \times 2.3$ mm³, 61 slices, 64 diffusion directions, b value=1000 s/mm², 1 b=0 image).

Structural and diffusion MRI processing

Detailed processing steps are covered in¹⁷¹⁸ with earlier iterations of the superficial white matter mapping methods covered in the work done by Phillips *et al.*⁸¹¹ The author responsible for MRI data analysis (OP) was blinded to clinical details of subjects. In brief, T1-weighted images were processed using BrainSuite's cortical surface extraction pipeline (http://brainsuite.org/processing/surfaceextraction/ v14b), which produces surface models of the cerebral cortex from T1 MRI.¹⁹ Next, the surfaces for each subject were registered to a reference atlas surface using BrainSuite's surface/volume registration software

Table 2 Sociodemographic and clinical characteristics of patients with anti-NMDAR encephalitis and healthy controls							
	Recovered patients (n=10)	Non-recovered patients (n=36)	Controls (n=30)	p Value			
Gender (man/woman)	2/8	4/32	6/24	0.471††, 1‡‡, 0.335§§			
Age (years) ±SD	26.6±7.38	26.7±8.83	27.5±8.42	0.359††, 0.546‡‡, 0.666§§			
mRS at MRI	0±0	1.75±0.12	NA	0.001††			
Maximal mRS*	3.8±0.3	4.2±0.2	NA	0.31††			
Verbal memory†	65.5±7.88	53.0±11.7	65.3±7.94	0.003††, 0.94‡‡,<0.001§§			
Visuospatial memory‡	28.9±5.65	24.18±8.00	27.8±5.12	0.091††, 0.58‡‡, 0.041§§			
Working memory§	-0.119±0.507	-0.174 ± 0.864	0.472±0.776	0.850††, 0.036‡‡, 0.007§§			
Attention¶	-0.304 ± 0.418	0.362±1.22	-0.371 ± 0.565	0.099††, 0.74‡‡, 0.006§§			
Executive function**	497.6±46.5	525.9±77.5	442.3±66.5	0.303††, 0.035‡‡,<0.001§§			

Mean values are reported with the SD.

*Maximal mRS=5, 21 patients; maximal mRS=4, 11 patients; maximal mRS=3, 13 patients; maximal mRS=2, 1 patient.

†RAVLT sum score; missing data for one non-recovered patient.

‡ROCF delayed recall; missing data for three non-recovered patients.

§Working memory composite score; missing data for one recovered patients and nine non-recovered patients.

¶Alertness score; missing data for two non-recovered patients.

**Go/No-Go reaction time; missing data for two non-recovered patients.

††Recovered patients versus non-recovered patients.

##Recovered patients vs controls.

§§Non-recovered patients vs controls.

Anti-NMDAR, anti-N-methyl-D-aspartate receptor; mRS, modified Rankin scale; NA, not available; RAVLT, Rey Auditory Verbal Learning Test; ROCF, Rey-Osterrieth Complex Figure.

Neuro-inflammation

v14b).^{20–22} (SVReg; http://brainsuite.org/processing/svreg/, Outputs from SVReg were inspected to ensure proper segmentation and surface/volume registration. The diffusion-weighted images were processed with the BrainSuite Diffusion Pipeline (BDP²³; http://brainsuite.org/processing/diffusion/). Finally, to allow cross-subject sampling of anatomically comparable superficial white matter mean diffusivity, diffusivity was sampled along each vertex of the white matter surface that had been mapped to the atlas reference via SVReg. Mean diffusivity was chosen to be the focus of this study rather than other diffusion metrics commonly examined in deep white matter investigations (axial and radial diffusivity, fractional anisotropy) given that the difference between axial and radial diffusivity values is considerably smaller in the superficial white matter than in the deep white matter.^{8 17 18} Methodological limitations include the inherent difficulty for tissue classification in MRI images and potential partial volume effects, however, these are minimised by the inspection of each tissue-classified image to ensure proper segmentation. Furthermore, we generated a ventricle/Cerebrospinal fluid (CSF) and subcortical grey matter mask (including the hippocampus which is not part of BrainSuite's surface model). This image mask was then applied to the diffusivity image to exclude these regions from analysis. This step was done as an additional quality control measure to maximise sampling of the superficial white matter and to minimise confounds from other areas.

Whole brain mean diffusivity was extracted for each subject and these values were then used to test for differences between groups (recovered patients, non-recovered patients and controls) using SPSS's General Linear Model (GLM) with sex and age as covariates. If a difference between groups was detected at the whole brain level, we followed up within lobar (frontal, temporal, parietal, limbic and occipital) regions of interest (ROIs) to identify where the changes were located. Likewise, for increased spatial resolution, we applied an exploratory GLM (http://brainsuite.org/bss/)²⁴ at the vertex level for lobar ROIs that were shown to be significantly different between groups to identify precisely where the differences in mean diffusivity were located.

Correlations between the superficial white matter and cognitive measures

To investigate whether superficial white matter changes were related to global degree of disability and cognitive scores (verbal memory (RAVLT sum score), visuospatial memory (ROCF delayed recall), attention (mean of z-transformed phasic and tonic alertness performance) and working memory (mean of z-transformed digit span backward and block span backward performance)), we ran a partial correlation analysis including all patients (recovered and non-recovered patients) between these measures and whole brain superficial white matter mean diffusivity with sex and age as covariates. If there was a significant effect, we investigated further within lobar ROIs. If there was a significant effect within an ROI we included this region in an exploratory vertex-based GLM analysis with sex and age as covariates between the measures and superficial white matter mean diffusivity. The significance threshold was set for all tests at p<0.05.

RESULTS

There was no significant difference between recovered and non-recovered patients regarding age (recovered, mean 26.60 ± 7.38 years vs non-recovered $26.69\pm8.83\pm1.8$ years;



Figure 1 Whole brain superficial white matter mean diffusivity in patients with anti-NMDAR encephalitis and healthy controls. * Indicates significant difference between groups. There was a significant difference between controls and non-recovered patients (p<0.001) but no significant difference between controls and recovered patients (p>0.05). Box plots show the average superficial white matter mean diffusivity value for each group. Mean diffusivity units: 10^{-3} mm²/s. NMDAR, anti-*N*-methyl-D-aspartate receptor.

p=0.471), CSF antibody titres (median 3.2 (range 0-10) vs 3.2 (0-320); p=0.84), maximal mRS scores (mean 3.8 ± 0.3 vs 4.2 ± 0.2 ; p=0.31), disease duration (duration of hospitalisation; mean 84.1±26.0 vs 105.6±18.8 days; p=0.55), time since disease onset (mean 23.9 ± 5.2 vs 26.5 ± 4.3 months; p=0.74), or time between symptom onset and first treatment (mean 224 ± 128 days vs 155 ± 40 days; p=0.65). Non-recovered patients had marked cognitive impairment and performed significantly worse than healthy controls in all investigated cognitive domains, that is, verbal memory, visuospatial memory, working memory, attention and executive function (table 2). In contrast, recovered patients had normal performance for visuospatial memory, verbal memory and attention and only mild deficits in working memory and attention. No correlation between antibody titres and superficial white matter changes or cognitive function wasobserved.

Whole brain analysis

Whole brain superficial white matter mean diffusivity analysis revealed a significant microstructural integrity impairment in non-recovered patients in comparison with recovered patients (recovered vs non-recovered, df=45, F=11.62, p=0.001) and healthy controls (non-recovered vs controls, df=64, F=13.098, p=0.001; figure 1). In contrast, no impairment of superficial white matter integrity was observed in recovered patients in comparison to controls (recovered vs controls, df=38, F=1.944, p=0.172).

Lobar region-of-interest analysis

To explore superficial white matter damage in non-recovered patients in more detail, we performed lobar ROI analyses that showed significant differences between patients and controls within the frontal, temporal and parietal lobes bilaterally as well as within the left occipital lobe (table 3; online supplementary figure 1).

 Table 3
 Whole brain and lobar superficial white matter mean diffusivity in non-recovered patients with anti-NMDAR encephalitis and healthy controls

	Controls		Non-recovered patients	df=64		
	Mean	SD	Mean	SD	F	p Value
Whole brain	9.18E-04	2.92E-05	9.53E-04	4.37E-05	13.098	0.001
Left frontal	9.12E-04	2.86E-05	9.57E-04	4.93E-05	10.408	0.002
Right frontal	9.14E-04	3.22E-05	9.62E-04	5.15E-05	12.649	0.001
Left temporal	8.50E-04	3.29E-05	8.95E-04	8.20E-05	16.822	0.001
Right temporal	8.75E-04	3.20E-05	9.09E-04	6.25E-05	17.079	0.001
Left parietal	9.52E-04	5.20E-05	9.68E-04	6.89E-05	6.838	0.011
Right parietal	9.57E-04	5.76E-05	9.80E-04	6.28E-05	7.766	0.007
Left limbic	1.00E-03	9.92E-05	1.07E-03	1.29E-04	0.958	0.332
Right limbic	1.04E-03	9.44E-05	1.06E-03	1.34E-04	3.057	0.085
Left occipital	9.36E-04	6.18E-05	9.54E-04	7.24E–05	5.601	0.021
Right occipital	9.26E-04	3.71E-05	9.37E-04	5.94E-05	0.321	0.573

Significant results are in bold.

NMDAR, anti-N-methyl-D-aspartate receptor.

Vertex-based analysis

To allow for a more fine-grained anatomical mapping, a precise vertex-based analysis was performed only within the lobar ROIs where there was a significant effect and revealed wide-spread differences (figure 2). In order to quantify which vertices had the largest differences in diffusivity between controls and non-recovered patients, we calculated the percentage difference at each vertex (figure 3). Percentage difference maps revealed increased mean diffusivity within the superficial white matter in distributed brain regions most prominently within the frontal, temporal and parietal lobes.

Correlation analysis within the superficial white matter

Whole brain superficial white matter damage was significantly correlated with disease severity (mRS) and cognitive deficits in verbal memory, visuospatial memory, attention and working memory, that is, a higher mean diffusivity was associated with a worse performance (table 4). In order to localise correlations in more detail, we performed lobar ROI analyses, which showed widespread associations in the same direction, that is, higher mean diffusivity was associated with a worse performance (table 4). In order to identify precisely which vertices were associated with clinical symptoms within significantly associated lobar ROIs only, we ran vertex-wise correlation analyses (figure 4). Verbal memory strongly correlated with superficial white matter damage throughout the brain, but particularly with damage in the temporal lobes bilaterally and the left cingulate, while visuospatial memory was associated with damage bilaterally in the lateral frontal lobes, the left temporal lobe and left cingulate. Working memory correlated with damage in the bilateral lateral



Figure 2 Superficial white matter damage in non-recovered patients with anti-NMDAR encephalitis. Probability maps show effects of NMDAR encephalitis on the superficial white matter (group difference between non-recovered patients and controls): mean diffusivity controlling for age and gender was mapped at high-spatial resolution at thousands of homologous locations within the superficial white matter. Vertex-based mapping was carried out only in lobar regions where there was a significant difference between non-recovered patients and controls. The direction of effects is indicated by the colour bar: purple/red/yellow colours indicate increased diffusivity and cyan/green/blue indicate reduced diffusivity in non-recovered patients relative to healthy controls. anti-NMDAR, anti-*N*-methyl-D-aspartate receptor.



Figure 3 Mean diffusivity percentage difference between non-recovered patients with anti-NMDAR encephalitis and healthy controls. Vertex based percentage difference maps show the difference between non-recovered patients and controls. Heat map colour bar indicates percentage difference between the two groups at each vertex point. Anti-NMDAR, anti-*N*-methyl-D-aspartate receptor.

|--|

	Dise	Disease severity		Verbal memory		Visuospatial memory		Attention		Working memory	
	r	p Value	r	p Value	r	p Value	r	p Value	r	p Value	
Whole brain	0.453	0.002	-0.597	0.001	-0.357	0.022	0.338	0.016	-0.347	0.044	
Left frontal	0.466	0.001	-0.563	0.001	-0.347	0.026	0.294	0.07	-0.472	0.001	
Right frontal	0.48	0.001	-0.464	0.002	-0.35	0.025	0.3	0.064	-0.503	0.001	
Left temporal	0.251	0.1	-0.336	0.028	-0.468	0.002	0.276	0.089	-0.215	0.223	
Right temporal	0.316	0.036	-0.491	0.001	-0.207	0.195	0.23	0.158	-0.212	0.23	
Left parietal	0.246	0.107	-0.441	0.003	-0.167	0.296	0.3	0.063	-0.032	0.063	
Right parietal	0.318	0.035	-0.547	0.001	-0.142	0.374	0.329	0.041	-0.329	0.856	
Left limbic	0.262	0.086	-0.405	0.007	-0.371	0.017	0.249	0.127	-0.149	0.402	
Right limbic	0.026	0.867	-0.045	0.774	0.138	0.39	-0.129	0.433	-0.129	0.433	
Left occipital	0.3	0.048	-0.473	0.001	-0.233	0.143	0.385	0.015	-0.144	0.417	
Right occipital	0.291	0.055	-0.389	0.01	-0.107	0.505	0.54	0.001	-0.145	0.414	

Pearson correlation coefficients and p values for the correlation between disease symptoms and white matter integrity in the whole brain and lobar regions of interest are shown. Correlations included all patients (recovered and non-recovered).

Results in bold font are statistically significant.

prefrontal regions. Attention correlated with damage predominantly in the occipital lobes bilaterally and in the right parietal lobe. Disease severity (mRS) was associated with superficial white matter damage in the frontal, temporal, parietal and occipital brain regions (figure 4).

DISCUSSION

The aim of this study was to investigate superficial white matter integrity in patients with anti-NMDAR encephalitis. The following main findings emerged from our investigation: (1) there are widespread abnormalities in the superficial white matter of non-recovered patients; (2) recovered patient's superficial white matter showed no abnormalities; (3) superficial white matter integrity is strongly correlated with disease symptoms and disease-related cognitive deficits. Previous brain imaging research on anti-NMDAR encephalitis has demonstrated abnormalities of the long-range deep white matter,² hippocampus³ and widespread alteration of functional connectivity with predominant affection of frontotemporal connections.² Our results are in line with these previous findings and additionally demonstrate that there are extensive abnormalities in brain structures that have not been examined previously. Importantly, these abnormalities are closely linked to cognitive impairments.

The observed results clearly indicate that abnormalities of the superficial white matter persist in non-recovered patients. However, the superficial white matter is a complex area and our knowledge of this region is still limited. For example, besides containing corticocortical short-range connections,¹¹ long-range axonal projections also pass through this area²⁵ and it additionally contains a high proportion of 'interstitial neurons'.²⁶ Volumetrically, the superficial white matter is largely composed of myelin and oligodendrocytes. Indeed, oligodendrocytes express NMDARs,^{27 28} that may result in a vulnerability to antibodies targeting these receptors and loss of myelin sheaths.²⁹ Together, this suggests that the observable diffusivity changes are driven by changes in oligodendrocytes. These changes could be particularly detrimental because oligodendrocytes in the superficial white matter myelinate up to 50 axon segments with fewer than 10 myelin membrane wraps.⁹ These late myelinating oligodendrocytes are thus structurally more complex and metabolically overextended compared with the deep white matter where oligodendrocytes tend to myelinate only a single axon segment with about 100 wraps.^{10 30} This suggests that damage to

oligodendrocytes in this region affects a greater number of axons segments than in the deep white matter.

The majority of patients with anti-NMDAR encephalitis recover well with prompt diagnosis and immunotherapy, although cognitive deficits frequently persist.¹⁶ Previous research has linked disease severity and disease duration to the extent of structural damage, for example, hippocampal atrophy and deep white matter damage, that in turn predicted the severity of persisting cognitive impairments.^{2 3} These findings are corroborated by the present study and extended to the superficial white matter: patients who had not fully recovered (mRS > 0) exhibited widespread damage of the superficial white matter, while, in contrast, recovered patients (mRS=0) showed no damage. Importantly, there were no differences in clinical parameters such as antibody titre, maximal mRS scores or disease duration between recovered and non-recovered patients. This shows that the heterogeneity in disease course and recoveryas assessed using the mRS-cannot be fully explained by these clinical variables. Possible further relevant factors might include differences in NMDAR antibody affinity or differences in the individual susceptibility to white matter damage, for example, susceptibility of oligodendrocytes to NMDAR dysfunction-mediated pathology. Damage of the superficial white matter was furthermore closely tied to cognitive impairments: diffusivity in the superficial white matter correlated with deficits in attention, working memory and verbal and visuospatial long-term memory. Overall, these results suggest that superficial white matter damage contributes to disease symptoms, but can be reversible in patients that recover. However, this conclusion will need to be confirmed in longitudinal studies with larger patient cohorts. Multimodal longitudinal studies will also help to further elucidate the contribution of different pathologies in anti-NMDAR encephalitis, for example, hippocampal atrophy, deep white matter damage and superficial white matter damage, to disease symptoms.

We categorised patients with a mRS score of 0 as *recovered-* nevertheless, recovered patients showed moderate deficits of working memory and executive function that are not reflected by the mRS and that might well interfere with everyday activities in university or work.¹⁶ These remaining cognitive deficits may indicate subtle superficial white matter changes and/or may indicate that different structural and functional disease mechanisms contribute to the cognitive deficits in anti-NMDAR encephalitis,



Figure 4 Correlation between superficial white matter damage and cognitive deficits in patients with anti-NMDAR encephalitis. All scatter plots reflect significant correlations between the cognitive domain and whole brain superficial white matter mean diffusivity. Vertex-based mapping was carried out only in lobar regions where there was a significant correlation between the cognitive domains in question (see table 3). For verbal memory (RAVLT sum score), visuospatial memory (ROCF delayed recall) and working memory (mean of z-transformed digit span backward and block span backward performance), higher scores indicate better performance; for attention (mean of z-transformed phasic and tonic alertness performance), higher scores indicate worse performance; for disease severity (mRS), higher scores indicate higher disease severity. In the vertex-based correlation maps, the direction of effects is indicated by the colour bar: purple/red/yellow colours indicate higher diffusivity with worse performance and cyan/green/blue indicate higher diffusivity with a better performance. Correlations included all patients (recovered and non-recovered). Anti-NMDAR, anti–N-methyl-D-aspartate receptor; mRS, modified Rankin scale.

Neuro-inflammation

likely including the previously observed correlation of memory deficits with impaired hippocampal functional connectivity,² disrupted functional connectivity within the medial temporal lobe network and hippocampal atrophy.³

Dysfunction of NMDARs closely links anti-NMDAR encephalitis to current concepts of schizophrenia pathophysiology.³ The glutamate hypothesis of schizophrenia posits that NMDAR hypofunction can induce a hyperdopaminergic state and can cause the major disease symptoms. Importantly, NMDAR antagonists such as phencyclidine and ketamine reliably induce positive and negative schizophrenia symptoms.^{31 32} Consequently, anti-NMDAR encephalitis and schizophrenia can have a remarkably similar clinical presentation with hallucinations, psychosis and delusions (positive symptoms), but also deficits of motivation, attention and memory (negative symptoms).^{33 34} In line with this pathophysiological and clinical overlap, there are notable similarities in neuroimaging findings between the two disorders, including reduced hippocampal volumes, abnormal resting state connectivity and deep white matter changes.^{2 3 5 35} Extending these observations, our current results in anti-NMDAR encephalitis mirror the impairments of the superficial white matter in schizophrenia.¹¹ For example, both non-recovered patients with anti-NMDAR encephalitis and patients with schizophrenia share widespread abnormalities, for example, in the temporal lobes bilaterally. Furthermore, vertex-based analyses show overlap between the two disorders in many focal locations including the lateral frontal lobes. The similarity of the results might indicate shared pathophysiological mechanisms that led to superficial white matter abnormalities in anti-NMDAR encephalitis and schizophrenia.

Importantly, anti-NMDAR encephalitis presents a unique opportunity to increase our understanding about brain function in general,³⁶ given that our findings illuminate the role of the superficial white matter in cognitive function. Verbal memory scores were linked with superficial white matter diffusivity across the whole brain, while visuospatial long-term memory was linked bilaterally with the frontal lobes as well as the left temporal lobe and cingulate gyrus. These observations are in line with recent evidence that challenges the view of a strict lateralisation of memory functions.³⁷ Working memory was closely tied to diffusivity in the frontal lobe bilaterally, corroborating extensive evidence from lesion, electrophysiological and functional MRI studies.^{38 39} Similarly, the observed correlation between visuospatial attention deficits and superficial white matter damage in the occipital lobes bilaterally and the right parietal lobe is in line with previous research.⁴⁰ This occipital damage might moreover contribute to the recently described visual dysfunction in patients with NMDAR encephalitis.¹⁵

CONCLUSION

The pathophysiological model of anti-NMDAR encephalitis, based on extensive immunocytochemical, physiological and molecular studies, holds that NMDAR antibodies induce a reversible capping and internalisation of NMDAR without complement activation and cell death.^{41 42} It might therefore be speculated that the widespread damage to the deep² and superficial white matter is the result of a downstream response to neuronal changes. Alternatively or in addition, interference of antibodies with NMDARs on oligodendrocytes might be involved in the widespread impairment of white matter in anti-NMDAR encephalitis. Independent of the exact mechanism still to be discovered, the role of white matter in the disorder seems to be more substantial than estimated previously.

Author affiliations

¹Department of Psychiatry, Division of Child and Adolescent Psychiatry, Stanford University School of Medicine, Stanford, California, USA

²Department of Neurology, Ahmanson Lovelace Brain Mapping Center, David Geffen School of Medicine at UCLA, Los Angeles, California, USA

³Department of Mental Health, King Faisal Specialist Hospital & Research Center, Riyadh, Saudi Arabia

⁴Départment of Clinical and Behavioral Neurology, IRCCS Santa Lucia Foundation, Rome, Italy

⁵Department of Neurology, Charité University Medicine Berlin, Berlin, Germany ⁶German Center for Neurodegenerative Diseases (DZNE), Berlin, Germany ⁷NeuroCure Clinical Research Center, Berlin, Germany

⁸Experimental and Clinical Research Center, Charité Universitätsmedizin, Berlin, Germany

⁹Experimental and Clinical Research Center, Max Delbrueck Center for Molecular Medicine, Berlin, Germany

¹⁰Berlin School of Mind and Brain, Humboldt-Universität zu Berlin, Berlin, Germany

Correction notice Since this paper was first published online, the author name Paul Friedmann has been updated to read Friedmann Paul.

Acknowledgements MS receives research support from the National Institute of Mental Health, the National Institute of Aging, the Office of Research in Women's Health, Brain and Behavior Research Foundation, Stanford Child Health Research Institute, Johnson & Johnson and Neuronetics, and she has served on an advisory board for Sunovion.

Contributors ORP and CF: wrote the manuscript and designed the analysis. SHJ, DWS and ORP: contributed to analysis methods development. CF, CJP, HP and FP: contributed to the acquisition, analysis and interpretation of data. All authors: contributed to the writing and editing of the manuscript and provided important intellectual content.

Funding This work was supported in part by the NIH National Institute of Neurological Disorders and Stroke award R01 NS074980 (DWS). FP is supported by Deutsche Forschungsgemeinschaft (DFG Exc 257).

Disclaimer The content is solely the responsibility of the authors and not an official position of the institution or funder and does not necessarily represent the official views of the National Institutes of Health.

Competing interests None declared.

Patient consent Obtained.

Ethics approval The study was approved by the Ethics Committee of the Charité - Universitätsmedizin Berlin.

Provenance and peer review Not commissioned; externally peer reviewed.

© Article author(s) (or their employer(s) unless otherwise stated in the text of the article) 2018. All rights reserved. No commercial use is permitted unless otherwise expressly granted.

REFERENCES

- Dalmau J, Lancaster E, Martinez-Hernandez E, et al. Clinical experience and laboratory investigations in patients with anti-NMDAR encephalitis. Lancet Neurol 2011;10:63–74.
- 2 Finke C, Kopp UA, Scheel M, et al. Functional and structural brain changes in anti-Nmethyl-D-aspartate receptor encephalitis. Ann Neurol 2013;74:n/a–96.
- 3 Finke C, Kopp UA, Pajkert A, et al. Structural Hippocampal Damage Following Anti-N-Methyl-D-Aspartate Receptor Encephalitis. Biol Psychiatry 2016;79:727–34.
- 4 Heine J, Ly LT, Lieker I, et al. Immunoadsorption or plasma exchange in the treatment of autoimmune encephalitis: a pilot study. J Neurol 2016;263:2395–402.
- 5 Peer M, Prüss H, Ben-Dayan I, et al. Functional connectivity of large-scale brain networks in patients with anti-NMDA receptor encephalitis: an observational study. Lancet Psychiatry 2017;4:768–74.
- 6 Dean DC, O'Muircheartaigh J, Dirks H, et al. Characterizing longitudinal white matter development during early childhood. Brain Struct Funct 2015;220:1921–33.
- 7 Bartzokis G. Age-related myelin breakdown: a developmental model of cognitive decline and Alzheimer's disease. *Neurobiol Aging* 2004;25:5–18.
- 8 Phillips OR, Clark KA, Luders E, et al. Superficial white matter: effects of age, sex, and hemisphere. Brain Connect 2013;3:146–59.
- 9 Butt AM, Berry M. Oligodendrocytes and the control of myelination in vivo: new insights from the rat anterior medullary velum. J Neurosci Res 2000;59:477–88.
- 10 Haroutunian V, Katsel P, Roussos P, et al. Myelination, oligodendrocytes, and serious mental illness. Glia 2014;62:1856–77.
- 11 Phillips OR, Nuechterlein KH, Asarnow RF, et al. Mapping corticocortical structural integrity in schizophrenia and effects of genetic liability. *Biol Psychiatry* 2011;70:680–9.
- 12 Titulaer MJ, McCracken L, Gabilondo I, *et al*. Treatment and prognostic factors for long-term outcome in patients with anti-NMDA receptor encephalitis: an observational cohort study. *Lancet Neurol* 2013;12:157–65.

- 13 Gresa-Arribas N, Titulaer MJ, Torrents A, et al. Antibody titres at diagnosis and during follow-up of anti-NMDA receptor encephalitis: a retrospective study. Lancet Neurol 2014;13:167–77.
- 14 Graus F, Titulaer MJ, Balu R, et al. A clinical approach to diagnosis of autoimmune encephalitis. Lancet Neurol 2016;15:391–404.
- 15 Brandt AU, Oberwahrenbrock T, Mikolajczak J, et al. Visual dysfunction, but not retinal thinning, following anti-NMDA receptor encephalitis. *Neurol Neuroimmunol Neuroinflamm* 2016;3:e198.
- 16 Finke C, Kopp UA, Prüss H, et al. Cognitive deficits following anti-NMDA receptor encephalitis. J Neurol Neurosurg Psychiatry 2012;83:195–8.
- 17 Phillips OR, Joshi SH, Piras F, et al. The superficial white matter in Alzheimer's disease. Hum Brain Mapp 2016;37:1321–34.
- 18 Phillips O, Joshi S, Squitieri F, et al. Major Superficial White Matter Abnormalities in Huntington disease. Frontiers 2016 http://journal.frontiersin.org/article/10.3389/fnins. 2016.00197/abstract
- 19 Shattuck DW, Leahy RM. BrainSuite: an automated cortical surface identification tool. Med Image Anal 2002;6:129–42.
- 20 Joshi AA, Shattuck DW, Thompson PM, et al. Surface-constrained volumetric brain registration using harmonic mappings. *IEEE Trans Med Imaging* 2007;26:1657–69.
- 21 Joshi AA, Shattuck DW, Thompson PM, et al. Cortical surface parameterization by p-harmonic energy minimization. Proc IEEE Int Symp Biomed Imaging 2004;1:428-431.
- 22 Joshi A, Shattuck DW LR. A fast and accurate method for automated cortical surface registration and labeling. *Fifth workship on biomedical image registration*. Nashville: Tennessee, 2012.
- 23 Bhushan C, Haldar JP, Choi S, *et al*. Co-registration and distortion correction of diffusion and anatomical images based on inverse contrast normalization. *Neuroimage* 2015;115:269–80.
- 24 Joshi SH, Woods RP, Joshi A, et al; A statistical toolbox for brainsuite: Organization for Human Brain Mapping (OHBM), 2014.
- 25 Reveley C, Seth AK, Pierpaoli C, et al. Superficial white matter fiber systems impede detection of long-range cortical connections in diffusion MR tractography. Proc Natl Acad Sci U S A 2015;112:E2820–E2828.
- 26 Suárez-Solá ML, González-Delgado FJ, Pueyo-Morlans M, et al. Neurons in the white matter of the adult human neocortex. Front Neuroanat 2009;3:7.
- 27 Káradóttir R, Cavelier P, Bergersen LH, et al. NMDA receptors are expressed in oligodendrocytes and activated in ischaemia. Nature 2005;438:1162–6.

- 28 Salter MG, Fern R. NMDA receptors are expressed in developing oligodendrocyte processes and mediate injury. *Nature* 2005;438:1167–71.
- 29 Paoletti P, Bellone C, Zhou Q. NMDA receptor subunit diversity: impact on receptor properties, synaptic plasticity and disease. *Nat Rev Neurosci* 2013;14:383–400.
- 30 Peters A, Sethares C. Oligodendrocytes, their progenitors and other neuroglial cells in the aging primate cerebral cortex. *Cereb Cortex* 2004;14:995–1007.
- 31 Masdeu JC, Dalmau J, Berman KF. NMDA Receptor Internalization by Autoantibodies: A Reversible Mechanism Underlying Psychosis? *Trends Neurosci* 2016;39:300–10.
- 32 Moghaddam B, Krystal JH. Capturing the angel in "angel dust": twenty years of translational neuroscience studies of NMDA receptor antagonists in animals and humans. *Schizophr Bull* 2012;38:942–9.
- 33 Dalmau J. NMDA receptor encephalitis and other antibody-mediated disorders of the synapse: The 2016 Cotzias Lecture. *Neurology* 2016;87:2471–82.
- 34 Lennox BR, Palmer-Cooper EC, Pollak T, et al. Prevalence and clinical characteristics of serum neuronal cell surface antibodies in first-episode psychosis: a case-control study. *Lancet Psychiatry* 2017;4:25–36.
- 35 Narr KL, Leaver AM. Connectome and schizophrenia. *Curr Opin Psychiatry* 2015;28:229–35.
- 36 De Bruijn M, Titulaer MJ. Chapter 12 Anti-NMDAR encephalitis and other glutamate and GABA receptor antibody encephalopathies. *Handbook of clinical neurology*, 2016:199–217.
- 37 Esfahani-Bayerl N, Finke C, Braun M, et al. Visuo-spatial memory deficits following medial temporal lobe damage: A comparison of three patient groups. *Neuropsychologia* 2016;81:168–79.
- 38 Curtis CE, D'Esposito M. Persistent activity in the prefrontal cortex during working memory. *Trends Cogn Sci* 2003;7:415–23.
- 39 Wager TD, Smith EE. Neuroimaging studies of working memory: a meta-analysis. Cogn Affect Behav Neurosci 2003;3:255–74.
- 40 Thiebaut de Schotten M, Dell'Acqua F, Forkel SJ, et al. A lateralized brain network for visuospatial attention. Nat Neurosci 2011;14:1245–6.
- 41 Hughes EG, Peng X, Gleichman AJ, et al. Cellular and synaptic mechanisms of anti-NMDA receptor encephalitis. J Neurosci 2010;30:5866–75.
- 42 Moscato EH, Peng X, Jain A, et al. Acute mechanisms underlying antibody effects in anti-N-methyl-D-aspartate receptor encephalitis. *Ann Neurol* 2014;76:108–19.